

Seabird Transfer of Nutrients and Trace Elements from the North Water Polynya to Land during the Mid-Holocene Warm Period, Carey Islands, Northwest Greenland

Peter M. Outridge,¹ Michael E. Goodsite,² Ole Bennike,³ Nicole Rausch⁴ and William Shotyk⁵

APPENDIX 1




FIG. S1. Interpolated dates from ¹⁴C age dating of Carey Islands peat core. (Regression equation: Calibrated date (yr BP) = $4.9 \times 10^3 + 16.1 * \text{Depth} - 2.72 \times 10^{-2} * (\text{Depth})^2$; $r^2 = 0.981$, $p < 0.001$. Dashed lines represent 95% confidence intervals.)

¹ Corresponding author: Centre for Earth Observation Science, University of Manitoba, 125 Dysart Road, Winnipeg, Manitoba R3T 2N2, Canada; Geological Survey of Canada, Natural Resources Canada, 601 Booth Street, Ottawa, Ontario K1A 0E8, Canada; peter.outridge@canada.ca

² Institute of Technology and Innovation, University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark

³ Geological Survey of Denmark and Greenland (GEUS), Øster Voldgade 10, DK-1350 Copenhagen K, Denmark

⁴ European Commission Joint Research Centre, Institute for Transuranium Elements, 76125 Karlsruhe, Germany

⁵ Department of Renewable Resources, University of Alberta, Edmonton, Alberta T6G 2H1, Canada