


Diversity and Keratin Degrading Ability of Fungi Isolated from Canadian Arctic Marine Bird Feathers

Brent M. Robicheau,^{1,2} Sarah J. Adams,¹ Jennifer F. Provencher,^{1,3} Gregory J. Robertson,⁴ Mark L. Mallory^{1,5}
and Allison K. Walker^{1,6}

(Received 6 February 2019; accepted in revised form 10 June 2019)

APPENDIX 1



Keratinase positive isolates. Darker vials indicate keratinolytic activity. Vials labeled 1322 are *Cadophora malorum*, those labeled 1321 are *Neosetophoma samarorum*, and those labeled 1319 are *Pseudogymnoascus pannorum*. In each image the vials to the right of a vial labeled Neg. CTRL are all negative controls where vials were not inoculated with fungi. *Pseudogymnoascus pannorum* was scored as a weak positive because only one of the three triplicates showed breakdown of the keratin azure.

¹ Department of Biology, Acadia University, 33 Westwood Ave., Wolfville, Nova Scotia B4P 2R6, Canada

² Present address: Department of Biology, Dalhousie University, 1355 Oxford St., Halifax, Nova Scotia B3H 4R2, Canada

³ Canadian Wildlife Service, Environment and Climate Change Canada, Gatineau, Québec K1A 0H3, Canada

⁴ Wildlife Research Division, Environment and Climate Change Canada, Mount Pearl, Newfoundland and Labrador A1N 4T3, Canada

⁵ Canada Fulbright Chair in Arctic Studies, University of Washington, Box 353650, Seattle, Washington 98195-3560, USA

⁶ Corresponding author: allison.walker@acadiau.ca