

Evaluating the outcomes of problem-based learning in postgraduate medical education: a systematic review and meta-analysis

Évaluation des résultats de l'apprentissage par problème dans la formation médicale postuniversitaire: une revue systématique et une méta-analyse

Emma A Mensour,^{1,2} Carolyn Tran,¹ Toni Li,³ Indika Mallawaarachchi,⁴ Jennifer M Shaw,² Sarah Blissett⁵

¹Schulich School of Medicine & Dentistry, Western University, Ontario, Canada; ²Centre for Education Research & Innovation (CERI), Schulich School of Medicine & Dentistry, Western University, Ontario, Canada; ³School of Medicine, Queen's University, Ontario, Canada; ⁴Department of Public Health Sciences, School of Medicine, University of Virginia, Virginia, USA; ⁵Department of Medicine, Division of Cardiology, London Health Sciences Center, Ontario, Canada

Correspondence to: Dr. Sarah Blissett, MD, MHPE, Schulich School of Medicine and Dentistry, Western University, 1151 Richmond Street, London, Ontario Canada N6A 5C1; email: sarah.blissett@lhsc.on.ca

Published ahead of issue: Jan 6, 2025; published: Feb 28, 2025. CMEJ 2025, 16(1). Available at <https://doi.org/10.36834/cmej.77394>

© 2025 Mensour, Tran, Li, Mallawaarachchi, Shaw, Blissett; licensee Synergies Partners. This is an Open Journal Systems article distributed under the terms of the Creative Commons Attribution License. (<https://creativecommons.org/licenses/by-nc-nd/4.0>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is cited.

Abstract

Background: Educators have recently been compelled to incorporate more active instructional formats into medical education, such as problem-based learning (PBL). In view of the mixed outcome data on the use of PBL in postgraduate medical education (PGME), there is a need to synthesize the data to inform the application of PBL in PGME contexts.

Objective: The aim of this systematic review and meta-analysis was to synthesize learning outcomes of PBL in PGME contexts.

Methods: The authors systematically searched MEDLINE, Embase, APA PsycINFO, AMED, CINAHL, Web of Science, ERIC, and Cochrane databases from January 1, 1950, to July 1, 2022 for original studies that reported Kirkpatrick outcomes of PBL in PGME contexts. Outcomes data were extracted. Quantitative data relating to learning outcomes were meta-analyzed using a random-effects model to generate weighted mean differences.

Results: Of 4310 abstracts screened, the authors included 21 studies encompassing anesthesia, family medicine, internal medicine, occupational medicine, pediatrics, psychiatry, public health and surgical residency programs. The studies reported reaction ($n = 12$), learning ($n = 15$), behavioural ($n = 6$) and/or results outcomes ($n = 4$). Meta-analysis of the three eligible articles demonstrated no significant difference after PBL in pre- and post-test results (pooled mean difference=0.13%, 95% CI, -6.74–7.00). There were observed improvements in satisfaction levels and self-reported behavioural outcomes following PBL.

Conclusions: Although similar learning outcomes were observed using PBL and the usual teaching in PGME, PBL was associated with benefits in trainee satisfaction and behavioural changes that contribute to learning and performance. PGME programs should consider incorporating PBL into curricula.

Résumé

Contexte : Les éducateurs doivent maintenant incorporer des formats pédagogiques plus actifs dans l'enseignement médical, tels que l'apprentissage par problèmes (APP). Compte tenu des données mitigées concernant l'utilisation l'APP dans la formation médicale postuniversitaire, il est nécessaire de synthétiser l'état des connaissances à cet effet, afin d'éclairer l'application de l'APP dans les contextes de formation médicale postuniversitaire.

Objectif : L'objectif de cette revue systématique et de cette méta-analyse était de synthétiser les travaux concernant l'APP effectués dans les contextes de formation médicale postdoctorale.

Méthodes : Les auteurs ont effectué des recherches systématiques dans les bases de données MEDLINE, Embase, APA PsycINFO, AMED, CINAHL, Web of Science, ERIC et Cochrane entre le 1^{er} janvier 1950 et le 1^{er} juillet 2022 pour identifier les études originales faisant état du niveau d'apprentissage basé sur le modèle de Kirkpatrick concernant l'APP dans des contextes de formation médicale postdoctorale. Les données relatives aux résultats ont été extraites. Les données quantitatives relatives au niveau d'apprentissage ont été méta-analysées à l'aide d'un modèle à effets aléatoires afin de générer des différences de moyennes pondérées.

Résultats : Sur les 4 310 résumés examinés, les auteurs ont inclus 21 études portant sur des programmes de résidence en anesthésie, en médecine familiale, en médecine interne, en médecine du travail, en pédiatrie, en psychiatrie, en santé publique et en chirurgie. Les études ont fait état des réactions ($n = 12$), de l'apprentissage ($n = 15$), du comportement ($n = 6$) et/ou des résultats ($n = 4$). La méta-analyse des trois articles admissibles n'a pas montré de différence significative après l'APP dans les résultats pré- et post-test (différence moyenne regroupée = 0,13 %, IC à 95 %, -6,74-7,00). Des améliorations ont été observées dans les niveaux de satisfaction et les résultats comportementaux autodéclarés à la suite de l'APP.

Conclusions : Bien que des résultats reliés à l'apprentissage similaires aient été observés en utilisant l'APP comparativement à l'enseignement habituel dans le cadre de la formation médicale postdoctorale, l'APP a été associé à des avantages en termes de satisfaction des apprenants et de changements de comportement qui contribuent à l'apprentissage et au rendement. Les programmes d'éducation médicale postdoctoraux devraient envisager d'intégrer l'APP dans les programmes d'études.

Introduction

Active instructional formats encourage students to engage, reflect, and contribute to the learning process. Compared with passive, teacher-centered instructional formats, active instructional formats have demonstrated improved knowledge retention and application.^{1,2} In recent years, medical education is shifting from passive to active instruction for its merits in improving cognition, collaboration, and self-directed learning skills, and intrinsic motivation.³

Problem Based learning (PBL) is an active instructional format. PBL is typically facilitated over multiple sessions. In the first session, a clinical problem is presented to small groups of students and led by a facilitator. Team members work together to understand the problem, identify questions to answer, solidify existing knowledge, acknowledge gaps in knowledge, and formulate learning objectives. After dedicated time for self-study (e.g., literature review, practical experience and/or formal lectures), the group reconvenes for a second session with the facilitator to discuss the findings and consolidate their learning. PBL has been incorporated across the spectrum of medical education, spanning from undergraduate medical education³ to continuing medical education.⁴

PGME trainees could benefit from PBL. In some studies in PGME contexts, but not all, there have been improved behavioural outcomes. PGME trainees were more competent in their communication, problem-solving, and self-learning skills, and showed more initiative than trainees instructed with passive learning instructional formats.⁵⁻⁷ While the effect of PBL on knowledge was variable—with studies favouring PBL⁸ or traditional lectures,⁹ and one showing no significant difference between the two groups¹⁰—students taught by PBL tended to outperform their conventional counterparts on clinical examinations and on faculty evaluations.^{11,12}

Yet, the application of PBL in PGME context has drawbacks. Transforming a teaching practice into PBL can be challenging as medical educators may have limited experiences with PBL structure and training. Faculty and teaching staff may be unfamiliar with PBL format and thus, tend to teach in the same passive instructional formats in which they were taught during their own undergraduate years.¹³ As a result, they may be slow to adapt the PBL format where needed.¹⁴ Another concern with implementing PBL in PGME is the material and educator resource investment. PBL instructional formats require

more time and potentially more facilitators to accommodate small group learning.¹⁵ Organizing sessions within the busy PGME context, with variable attendance due to vacations and post-call days, also represents a challenge. Level of knowledge acquisition,^{16,17} facilitator buy-in,¹⁵ and scheduling logistics in PBL are additional barriers to implementing PBL in PGME contexts.

In view of the mixed study outcomes data on the use of PBL in PGME contexts, training programs may be reluctant to embrace PBL. There is a need to consolidate and synthesize the data to determine the application of PBL formats across all areas of PGME to guide whether implementing PBL should be undertaken by PGME programs. Accordingly, the purpose of this systematic review and meta-analysis is to synthesize the available literature and make sense of conflicting results of PBL studies in PGME.

Methods

Selection of studies

In 2022, the authors performed a systematic search of MEDLINE, Embase, APA PsycINFO, AMED, CINAHL, Web of Science, ERIC, and Cochrane from January 1, 1950 to July 1, 2022. The following subject headings were used to search for studies: “Problem-based Learning,” “Education,” “Medical,” and “Internship and Residency”. For non-subject heading-based databases, the authors search strategy was as follows: (“Problem-Based Learning”) AND (“graduate medical education”) OR (“Internship”) OR (“Residency”).

The authors managed the articles using the web-based software Covidence.¹⁸ Duplicate articles were removed, and two authors independently performed title and abstract screening (CT and JS) to identify the articles for full text review. Conflicts were resolved collaboratively or with a third investigator if uncertainties persisted. Full text review was subsequently performed by one author (CT). This paper followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) reporting guidelines.¹⁹

Eligibility criteria

The authors included original studies of PBL in PGME contexts that reported learner outcomes described by the Kirkpatrick Evaluation Model for analysis. The model comprises four levels of criterion: “reaction” assesses participant satisfaction and perception of the program, often through qualitative scales or structured interviews; “learning” measures the effect of training on knowledge

and skill as evaluated by scored examinations pre- and post-training; “behaviour” describes an observable shift in participant thinking or mindset that leads to practical changes; and “results” specifies how these changes impact practice or patient care. The authors excluded articles that were in an undergraduate medical education context, described outcomes outside the Kirkpatrick criteria, were published in a non-English language, or published as an abstract, letter, or conference paper.

Data extraction and analysis

Each of the two reviewers (CL and TL) independently extracted the country of publication, medical specialty of participants, PBL format, and quantitative and qualitative outcomes data. The reviewers independently appraised the study quality using the Medical Education Research Study Quality Instrument (MERSQI), which evaluates 10 criteria in six domains.²⁰ This tool evaluates on a 3-point scale the design, sampling, data type, validity of evaluation instruments, data analysis, and outcomes of a given study for a maximum score of 18.

Data relating to learning outcomes were meta-analyzed using a random-effects model to generate weighted mean differences using Stata 17.²¹ The authors chose to perform the analysis in two steps, first assuming a higher correlation estimate of 0.8 and then assuming no correlation. A *p*-value of <0.05 was considered statistically significant.

Author positionality

The authors represented medical students (EM, CT, TL), a research assistant (JS), a statistician (IM), and a program director of a PGME training program (SB). At the outset of the study, the authors held shared beliefs about the value of active learning and were open-minded about how PBL could impact Kirkpatrick outcomes.

Results

The literature search returned 4310 unique studies. (Figure 1) After title and abstract screening, the authors reviewed 97 full-text articles. Of these, the authors excluded 61 articles due to inappropriate outcomes (*n* = 33), methods (*n* = 12), or population (*n* = 9); non-English language (*n* = 5); or duplicate articles missed by Covidence (*n* = 2). Twenty-one studies were included for analysis.




Figure 1. Prisma Chart

Study characteristics

An overview of study characteristics is summarized in Table 1 and Table 2. The studies were published between 1989–2018, with the majority conducted in the United States (*n* = 13). There was a diverse background of specialties including anesthesia, family medicine, internal medicine, occupational medicine, pediatrics, psychiatry, public health and surgery. Most studies were observational (*n* = 13). The authors appraised the study quality using the MERSQI tool and is summarized in Table 3. The mean score was 11.07 +/- 3.38, with four studies having scores less than nine. PBL sessions were either embedded into existing academic training program (*n* = 13) or were set as an additional session outside of the program time (*n* = 8). PBL sessions were facilitated in person (*n* = 19), virtually (*n* = 1), or both virtually and in person instruction (*n* = 1).

Kirkpatrick’s outcomes

The Kirkpatrick model evaluates training programs in their ability to meet the needs of the organization and participants through measuring participant reaction, learning outcomes, behavioural changes, and results.²² As summarized in Table 1, most studies reported reaction (*n* = 12) and learning outcomes (*n* = 15), with fewer reporting behavioural (*n* = 6) or results (*n* = 4) outcomes. Nine studies reported more than one Kirkpatrick outcome.

Table 1.

| Resident Speciality | Author(s) and Year | Reference number | Country | Number of Participants (intervention; control) | Virtual or In person PBL | Duration of PBL | Measurement Tool(s) for PBL | Kirkpatrick Scale level Evaluated | | | |
|-----------------------|------------------------|------------------|----------------|--|--------------------------|--|---|-----------------------------------|----------|-----------|---------|
| | | | | | | | | Reaction | Learning | Behaviour | Results |
| Anesthesia | Carrero (2007) | 37 | Spain | 54 (29;25) | In-person | 90 min PBL session daily for two days for two consecutive years | Pre/Post PBL tests scores | 0 | 1 | 0 | 0 |
| | DeOliveiraFilho (2005) | 40 | Brazil | 12 (5;7) | In-person | Two 60 min PBL sessions for two academic terms | Pre/Post PBL tests scores | 0 | 1 | 0 | 0 |
| | Komasawa (2018) | 38 | United States | 35 (35;0) | In person | Six to eight hour PBL sessions for one day for three consecutive years | Self confidence survey for anesthesia related crisis management | 0 | 0 | 1 | 0 |
| | Sakai (2013) | 41 | United States | 178 (93;85) | In person | 90 min PBL for four consecutive years. | Consultation solicitations & research projects | 0 | 0 | 1 | 0 |
| Family Medicine | Grol (1989) | 39 | Netherlands | 63 (31;32) | In person | Weekly PBL sessions for one year | Maastricht progress test | 0 | 1 | 1 | 1 |
| | Prislin (1997) | 34 | United States | 85 (27;58) | In person | Weekly PBL sessions for six years | Pre/post PBL 5-point Likert surveys | 0 | 1 | 0 | 1 |
| | Sass (2001) | 24 | United States | 9(9;0) | In person | Weekly PBL sessions for four weeks | Post-rotation Interview | 1 | 1 | 1 | 0 |
| Internal Medicine | Cook (2009) | 25 | United States | 105 (55, 50)* | Virtual | Weekly 45 min PBL sessions for one year | Pre/Post PBL tests scores | 1 | 1 | 0 | 0 |
| | Foley (1993) | 26 | United States | 7(7;0) | In person | 34 hours of PBL sessions per week for two months | Post-rotation Interview | 1 | 0 | 0 | 0 |
| | Mulder (2009) | 36 | Netherlands | 91(91;0) | In person | Once weekly two hour PBL sessions for eight weeks | Pre/Post PBL tests scores | 0 | 1 | 0 | 0 |
| Occupational medicine | Smits (2003) | 9 | Netherlands | 118 (59;59) | In person | Once weekly one day PBL session for four weeks | Pre/post PBL test scores and performance review | 1 | 1 | 0 | 1 |
| Pediatrics | Goodyear (2005) | 27 | United Kingdom | 14 (7;7) | In person | Not provided | OSCE and Pre/Post course self assessment sheets | 1 | 1 | 0 | 1 |
| | Ozuah (2001) | 28 | United States | 80 (39;41) | In person | Twice weekly PBL sessions for three months | Self administered questionnaire | 1 | 0 | 1 | 0 |
| Psychiatry | Heru (2011) | 29 | United States | 9 (9;0) | In person | Eight PBL Sessions over three weeks | Informal Testimony | 1 | 0 | 0 | 0 |
| | Schultz-Ross (1999) | 30 | United States | 23 (23;0) | In person | Three PBL sessions per year for 2 consecutive years | 5-point Likert-type Scales | 1 | 1 | 0 | 0 |
| | Shefet (2011) | 31 | Israel | 93 (93;0) | In person | Two-day PBL sessions | Satisfaction feedback sheets | 1 | 1 | 1 | 0 |
| | Yates (1996) | 33 | United States | 12 (12;0) | In person | Weekly PBL sessions for one year | Educational experiences survey | 1 | 0 | 0 | 0 |
| Public Health | Heading (2007) | 23 | Australia | 9 (9;0) | In person | Three PBL sessions over four days | Semi-structured interviews | 1 | 1 | 0 | 0 |
| Surgical | Itani (1997) | 43 | United States | 64 (64;0) | In person | Once weekly 60 min PBL session for one year | ABSITE questionnaire | 0 | 1 | 0 | 0 |
| | Lee (2008) | 35 | United States | 42 (42;0) | Virtual and in person | Weekly PBL sessions for two consecutive years | ABSITE questionnaire and scores | 0 | 1 | 0 | 0 |
| | Nguyen (2006) | 32 | United States | 55 (55;0) | In person | Weekly PBL sessions for six months | ABSITE questionnaire and scores, resident satisfaction survey | 1 | 1 | 0 | 0 |
| Total | | | | | | | | 12 | 15 | 6 | 4 |

Table 2

| Resident Speciality | Author(s) and Year | Reference number | Country | Number of Participants (intervention; control) | Virtual or In-person PBL | Duration of PBL | Measurement Tool(s) for PBL | Kirkpatrick Scale level Evaluated | | | |
|-----------------------|------------------------|------------------|----------------|--|--------------------------|--|---|-----------------------------------|----------|-----------|---------|
| | | | | | | | | Reaction | Learning | Behaviour | Results |
| Anesthesia | Carrero (2007) | 37 | Spain | 54 (29;25) | In-person | 90 min PBL session daily for two days for two consecutive years | Pre/Post PBL tests scores | 0 | 1 | 0 | 0 |
| | DeOliveiraFilho (2005) | 40 | Brazil | 12 (5;7) | In-person | Two 60 min PBL sessions for two academic terms | Pre/Post PBL tests scores | 0 | 1 | 0 | 0 |
| | Komasawa (2018) | 38 | United States | 35 (35;0) | In person | Six to eight hour PBL sessions for one day for three consecutive years | Self confidence survey for anesthesia related crisis management | 0 | 0 | 1 | 0 |
| | Sakai (2013) | 41 | United States | 178 (93;85) | In person | 90 min PBL for four consecutive years. | Consultation solicitations & research projects | 0 | 0 | 1 | 0 |
| Family Medicine | Grol (1989) | 39 | Netherlands | 63 (31;32) | In person | Weekly PBL sessions for one year | Maastricht progress test | 0 | 1 | 1 | 1 |
| | Prislin (1997) | 34 | United States | 85 (27;58) | In person | Weekly PBL sessions for six years | Pre/post PBL 5-point Likert surveys | 0 | 1 | 0 | 1 |
| | Sass (2001) | 24 | United States | 9(9;0) | In person | Weekly PBL sessions for four weeks | Post-rotation Interview | 1 | 1 | 1 | 0 |
| Internal Medicine | Cook (2009) | 25 | United States | 105 (55, 50)* | Virtual | Weekly 45 min PBL sessions for one year | Pre/Post PBL tests scores | 1 | 1 | 0 | 0 |
| | Foley (1993) | 26 | United States | 7(7;0) | In person | 34 hours of PBL sessions per week for two months | Post-rotation Interview | 1 | 0 | 0 | 0 |
| | Mulder (2009) | 36 | Netherlands | 91(91;0) | In person | Once weekly two hour PBL sessions for eight weeks | Pre/Post PBL tests scores | 0 | 1 | 0 | 0 |
| Occupational medicine | Smits (2003) | 9 | Netherlands | 118 (59;59) | In person | Once weekly one day PBL session for four weeks | Pre/post PBL test scores and performance review | 1 | 1 | 0 | 1 |
| Pediatrics | Goodyear (2005) | 27 | United Kingdom | 14 (7;7) | In person | Not provided | OSCE and Pre/Post course self assessment sheets | 1 | 1 | 0 | 1 |
| | Ozuah (2001) | 28 | United States | 80 (39;41) | In person | Twice weekly PBL sessions for three months | Self administered questionnaire | 1 | 0 | 1 | 0 |
| Psychiatry | Heru (2011) | 29 | United States | 9 (9;0) | In person | Eight PBL Sessions over three weeks | Informal Testimony | 1 | 0 | 0 | 0 |
| | Schultz-Ross (1999) | 30 | United States | 23 (23;0) | In person | Three PBL sessions per year for 2 consecutive years | 5-point Likert-type Scales | 1 | 1 | 0 | 0 |
| | Shefet (2011) | 31 | Israel | 93 (93;0) | In person | Two-day PBL sessions | Satisfaction feedback sheets | 1 | 1 | 1 | 0 |
| | Yates (1996) | 33 | United States | 12 (12;0) | In person | Weekly PBL sessions for one year | Educational experiences survey | 1 | 0 | 0 | 0 |
| Public Health | Heading (2007) | 23 | Australia | 9 (9;0) | In person | Three PBL sessions over four days | Semi-structured interviews | 1 | 1 | 0 | 0 |
| Surgical | Itani (1997) | 43 | United States | 64 (64;0) | In person | Once weekly 60 min PBL session for one year | ABSITE questionnaire | 0 | 1 | 0 | 0 |
| | Lee (2008) | 35 | United States | 42 (42;0) | Virtual and in person | Weekly PBL sessions for two consecutive years | ABSITE questionnaire and scores | 0 | 1 | 0 | 0 |
| | Nguyen (2006) | 32 | United States | 55 (55;0) | In person | Weekly PBL sessions for six months | ABSITE questionnaire and scores, resident satisfaction survey | 1 | 1 | 0 | 0 |
| Total | | | | | | | | 12 | 15 | 6 | 4 |

Reaction and satisfaction outcomes

The studies used participant surveys, feedback sheets, interviews, and testimonies to assess participant reaction to the PBL intervention.^{9,23–33} Eight of 12 studies concluded high satisfaction, favourable attitudes, and preference for PBL over traditional lectures.^{24,26,28–33} (Tables 1 and 2) Of the two studies that did not improve attitudes or satisfaction in learners after PBL, one group cited appreciation for the variety of PBL offered despite there being no significant difference in satisfaction.²⁷ The PBL group in Smits et al. was the only included study to report lower satisfaction as compared to the lecture-based group in post-intervention and follow-up interviews.⁹ The authors hypothesized that the lower satisfaction was due to familiarity with passive formats and insufficient orientation to active formats.

Learning and knowledge outcomes

Knowledge acquisition was assessed using test scores, including American board exam scores in surgery and family practice^{9,24,25,27,30–32,34–40} and scores on Objective Structured Clinical Examinations.²⁷ Of the 15 studies that assess learning and knowledge outcomes, four studies had control groups comparing quantitative data.^{9,27,39,40} Pre- and post-intervention quantitative data were available for six of 15 studies.^{9,35–37,39,40} Other studies collected qualitative data through learner perception of knowledge or did not report test scores.^{24,26,30,31,33,34,38,39,41} Meta-analysis of only those that included control groups and reported pre and post intervention scores comprised three studies^{9,37,40} and demonstrated no significant difference

after intervention in pre- and post-test results between PBL and the instructional format in the control group (pooled mean difference= 0.13%, 95% CI, -6.74–7.00). (Figure 2) The PBL sessions of the three studies included in the meta-analysis followed a typical PBL format as previously described.⁴²

Behaviour outcomes

Behaviours studied included appreciation for the social determinants of health,²⁴ confidence in using demographic data,²⁴ confidence in management of anaesthesia-related crises,⁴¹ specific resource consultation in research,⁴¹ research ability,⁴¹ prevalence of self-directed learning,²⁸ and teamwork.³¹ All behaviours were self-reported and showed an improvement in all domains. Of note, self-directed learning was rated higher with PBL, though there was no significant long-term effect.²⁸

Results outcomes

While direct patient outcomes were not measured in the results, clinical skills were assessed by expert evaluation of consultation skills and medical performance, clinical review using performance indicators and Likert surveys in behavioural science practice activity. The effects of PBL on results were mixed: PBL was associated with significantly higher performance indicator scores compared with lecture-based learning;⁹ there was a significant difference in provided care in some behavioural conditions after PBL,³⁴ and a PBL group had improved medical performance while the traditional learning group showed better consultation skills.³⁹




Figure 2. Forest plot depicting meta-analysis of learning and knowledge outcomes

Table 3.

| Author (year) | Reference number | Study design* (number of Institutions) | Response Rate, % \bar{x} | Type of Data § | Validity of Evaluation Instruments or measures | | | Data Analyses¶ | | Highest Level of Outcome assessed | MERSQI score |
|------------------------|------------------|--|----------------------------|--|--|------------------|--------------------|-----------------|-----------------------------|-----------------------------------|--------------|
| | | | | | Internal Structure | Content Validity | Criterion validity | Appropriateness | Sophistication | | |
| Carrero (2007) | 37 | RCT (2) | ≥75% | Objective measure | NR | Reported | Reported | Appropriate | Beyond descriptive analysis | Knowledge | 15 |
| Cook (2009) | 25 | RCT (1) | ≥75% | Objective measure | Reported | Reported | Reported | Appropriate | Beyond descriptive analysis | Knowledge | 15.5 |
| deOliveiraFilho (2005) | 40 | Single group pre/post test(1) | NA | Objective measure | Reported | Reported | NR | Appropriate | Beyond descriptive analysis | Knowledge | 11.5 |
| Foley (1993) | 26 | Single group cross sectional (2) | NA | Subjective assessment by study subject | NA | NA | NA | Appropriate | Comparisons made | Satisfaction | 6 |
| Goodyear (2005) | 27 | Nonrandomized 2 groups (1) | ≥75% | Subjective assessment by study subject | Reported | Reported | Reported | Appropriate | Beyond descriptive analysis | Results | 12 |
| Grol (1989) | 39 | Nonrandomized 2 groups (1) | ≥75% | Objective measure | Reported | Reported | Reported | Appropriate | Beyond descriptive analysis | Results | 14.5 |
| Heading (2007) | 23 | Single group cross sectional (3) | NA | Subjective assessment by study subject | Reported | NR | Reported | Appropriate | Comparisons made | Knowledge | 8.5 |
| Heru (2011) | 29 | Single group post test only (1) | NA | Subjective assessment by study subject | NA | NA | NA | Appropriate | Comparisons made | Satisfaction | 5.5 |
| Itani (1997) | 43 | Single group post test only (1) | NA | Objective measure | NA | NA | NA | Appropriate | Beyond descriptive analysis | Knowledge | 10 |
| Komassawa (2018) | 38 | Single group cross sectional (2) | NA | Objective measure | Reported | Reported | Reported | Appropriate | Descriptive analysis only | Behaviour | 11.5 |
| Lee (2008) | 35 | Nonrandomized 2 groups (1) | NA | Objective measure | Reported | Reported | Reported | Appropriate | Beyond descriptive analysis | Knowledge | 13 |
| Mulder (2009) | 36 | Single group pre/post test(1) | <50% | Subjective assessment by study subject | Reported | Reported | Reported | Appropriate | Beyond descriptive analysis | Knowledge | 10.5 |
| Nguyen (2006) | 32 | 2 group nonrandomized (1) | NA | objective measure | Reported | Reported | Reported | Appropriate | Beyond descriptive analysis | Knowledge | 13 |
| Ozuah (2001) | 28 | 2 group nonrandomized (1) | ≥75% | Subjective assessment by study subject | NR | NR | Reported | Appropriate | Beyond descriptive analysis | Behaviour | 10.5 |
| Prislin (1997) | 34 | 2 group nonrandomized (1) | 50-74% | Objective measure | Reported | NR | Reported | Appropriate | Beyond descriptive analysis | Results | 13 |
| Sakai (2013) | 41 | 2 group nonrandomized (1) | 50-74% | Objective measure | NR | NR | NR | Appropriate | Beyond descriptive analysis | Behaviour | 11 |
| Sass (2001) | 24 | Single group post test only (1) | NA | Subjective assessment by study subject | NA | NA | NA | Appropriate | Comparisons made | Behaviour | 5 |
| Schultz-Ross (1999) | 30 | Single group pre/post test(1) | NA | Subjective assessment by study subject | Reported | NR | reported | Appropriate | Beyond descriptive analysis | Knowledge | 9.5 |
| Shefet (2011) | 31 | Single group post test only (1) | <50% | Subjective assessment by study subject | NA | NA | NA | Appropriate | Comparisons made | Behaviour | 6 |
| Smits (2003) | 9 | RCT (1) | ≥75% | Objective measure | Reported | Reported | Reported | Appropriate | Beyond descriptive analysis | Results | 15.5 |
| Yates (1996) | 33 | RCT (1) | ≥75% | Objective measure | Reported | Reported | Reported | Appropriate | Beyond descriptive analysis | Satisfaction | 15.5 |

Discussion

This systematic review and meta-analysis synthesizes the current body of knowledge on PBL outcomes in PGME context. Our results highlight how PBL had similar learning outcomes to traditional teaching formats in PGME contexts, with the added benefits of increased satisfaction and facilitation of behavioural changes.

While critics of PBL in PGME highlight logistical issues, PBL was logistically feasible in a multitude of PGME contexts. PBL was successfully integrated into one-day workshop^{37,38} and over longer time periods, such as within full-year programs.^{26,33,34,36,39,43} The flexibility of the PBL structure allowed programs to achieve their specific objectives in the constraints of the local environment while also maintaining PBL principles.

PBL increases resident satisfaction. While this may seem less important than other outcomes, learner satisfaction is an important factor for effective and continuous improvement of PGME programs.^{44,45} Previous reviews surveying PBL in UGME also highlight how PBL improves satisfaction: PBL was reported by medical students as “nurturing and enjoyable.”¹¹ The high level of satisfaction and preference of PBL over traditional lectures is relevant for educators as it reflects the positive perception of PBL.^{38,39} The lower satisfaction with PBL due to learner expectations as reported by Smits et al. highlights the importance of orienting learners to the expectations and active nature of PBL.⁹

Additionally, PBL increases behaviours beyond what was achieved with usual formats. The increase in positive learner behaviour, including self-directed learning,²⁸ research,⁴¹ analysis of clinical problems,³¹ and confidence⁹ after PBL in PGME contexts is an important finding for educators. Engaging regularly in self-directed learning, collaboration, and confidence-building allows physicians to provide the most informed care, contribute to medical knowledge, and create a cycle of growth whereby increasingly challenging presentations offer continuing opportunities to learn.

Interestingly, PBL may not improve learning outcomes as currently measured. The PBL approach places less emphasis on achieving high assessment scores, so learners are more likely to study for understanding rather than exams.^{7,17} It is important to recognize that of 15 studies reporting learning outcomes in this review, a minority of studies had control groups comparing quantitative data on PBL to traditional formats.^{9,27,39,40} It is also difficult to

control for confounding factors such as instructor variability or learner availability in the context of work and on-call demands.^{7,45}

The impact of PBL on result outcomes were mixed; few studies evaluated patient outcomes and some indirectly evaluated clinical performance. While result outcomes are often considered to be the most important, Cook et al.⁴⁶ highlight several challenges of focusing on patient outcomes including potentially biased outcome selection and failure to establish a causal link. These perspectives highlight that results outcomes may not be feasible. In our study, these challenges were exemplified in four studies that indirectly assessed supervisors’ evaluation of learner performance with no standardized rating system and were subject to biased outcome selection.^{9,34,39,40} Thus, results outcomes may not always be possible to study in PGME contexts.⁴⁶

Similar findings have been reported with PBL in undergraduate medical education contexts. A recent scoping review of 124 studies⁶ identified that undergraduate medical learners were highly satisfied with PBL, and PBL was more effective than other formats in improving problem solving skills and behaviors such as communication skills and self-learning skills. Like the PGME context, there were mixed knowledge outcomes however PBL was consistently no worse than traditional formats for academic performance.⁶ Logistical issues and greater human resources are also barriers in undergraduate medical education contexts.⁶

There are several gaps in the literature regarding PBL in PGME that require further investigation. Firstly, in a CBME era, future studies could explore PBL on growth mindset. Fostering a growth mindset necessitates active engagement, effort, and focus;⁴⁷ PBL offers this by scheduling time for collaborative exploration, self-study, and facilitated discussion with an expert of presented cases. While no studies formally assessed the mindset of learners, the goals of PBL on skills development and motivation are in alignment with that of a growth mindset and is seen in the effects on behaviour and reaction.⁴⁸ Secondly, few studies evaluated patient outcomes, possibly due to the difficulty in isolating one learner’s contributions in collaborative care. With initiatives such as the GEMINI database, Canada’s largest hospital data and analytics network,⁴⁹ future studies may be able to link teaching interventions with patient outcomes. Lastly, while there is emerging research examining the delivery of PBL in an online environment,^{25,50} literature reporting on virtual

delivery of PBL in PGME is scarce. In a post-COVID-19 pandemic era, future studies should investigate the option of virtual delivery of PBL and the use of learning technologies within PBL formats.

There are limitations that merit discussion. Firstly, “problem-based learning” is not a rigidly defined approach and is similar to other forms of active learning such as case-based learning, forms of small group learning, and flipped classroom models. PBL was chosen as it was specific to medical education, though the term’s ambiguity may have resulted in some studies being excluded in this review. Secondly, while frequency and duration of PBL formats were reported for each article, other PBL design variations and facilitator differences were not reported in the studies limiting the ability for the authors to further evaluate these PBL models. Thirdly, only 45% ($n = 9$) of studies included a control group, few studies captured all the PBL outcomes of interest, and few studies captured long term patient and resident outcomes of PBL in PGME. Fourthly, across the 21 studies included, the mean MERSQI score was 11.07 +/- 3.38 (range of 5-15.5, of possible 18). The reliance on self-reporting questionnaires also represents important limitations. To overcome these limitations, future studies should aim to include a report of all PBL design and facilitation details as well as outcomes, with a randomized control trial design.

Conclusion

PBL in PGME has its merits in strengthening outcomes of education including attitudes and behaviours of residents at no detriment to their acquisition of medical knowledge. While there may be initial reservations in implementing PBL, existing infrastructure used for teaching residents can facilitate the scheduling and transition to PBL in a multitude of PGME contexts. The implementation of PBL can be a valuable educational strategy for residents and medical educators in PGME programs.

Conflicts of Interest: The authors declare they have no conflicts of interest. The authors alone are responsible for the content and writing of the article.

Funding: There are no funding sources to report.

Edited by: Eric Heinz (section editor); Marco Zaccagnini (senior section editor); Marcel D’Eon (editor-in-chief)

Authorship: Mensour and Tran contributed equally to this work and share first authorship

References

1. Prober CG, Khan S. Medical education reimaged: A call to action. *Acad Med*. 2013;88(10):1407–10. <https://doi.org/10.1097/ACM.0b013e3182a368bd>
2. Prober CG, Heath C. Lecture halls without lectures — a proposal for medical education. *N Engl J Med*. 2012 May 3;366(18):1657–9. <https://doi.org/10.1056/NEJMp1202451>
3. Trullàs JC, Blay C, Sarri E, Pujol R. Effectiveness of problem-based learning methodology in undergraduate medical education: a scoping review. *BMC Med Ed* 2022 22:1. 2022 Feb 17;22(1):1–12. <https://doi.org/10.1186/s12909-022-03154-8>
4. Al-azri H, Ratnapalan S. Problem-based learning in continuing medical education L’ apprentissage par problèmes en éducation médicale continue. *Can Fam Phys*. 2014;60:157–65.
5. Donner RS, Bickley H. Problem-based learning: an assessment of its feasibility and cost. *Hum Pathol*. 1990;21(9):881–5. [https://doi.org/10.1016/0046-8177\(90\)90170-A](https://doi.org/10.1016/0046-8177(90)90170-A)
6. Trullàs JC, Blay C, Sarri E, Pujol R. Effectiveness of problem-based learning methodology in undergraduate medical education: a scoping review. *BMC Med Educ*. 2022;22(1):1–12. <https://doi.org/10.1186/s12909-022-03154-8>
7. Albanese MA, Mitchell S. Problem-based learning: a review of literature on its outcomes and implementation issues. *Acad Med* 1993 Jan;68(1):52–81. <https://doi.org/10.1097/00001888-199301000-00012>.
8. Suliman S, Al-Mohammed A, Al Mohanadi D, et al. It is all about patients’ stories: case-based learning in residents’ education. *Qatar Med J*. 2019:17. <http://dx.doi.org/10.5339/qmj.2019.17>
9. Smits PB, de Buissonjé CD, Verbeek JH, van Dijk FJ, Metz JC, ten Cate OJ. Problem-based learning versus lecture-based learning in postgraduate medical education. *Scand J Work Environ Health*. 2003;29(4):280–7. <https://doi.org/10.5271/sjweh.732>
10. Santos-Gomez L, Kalishman S, Rezler A, Skipper B, Mennin SP. Residency performance of graduates from a problem-based and a conventional curriculum. *Med Educ*. 1990 Jul 1;24(4):366–75. <https://doi.org/10.1111/j.1365-2923.1990.tb02453.x>.
11. Albanese MA, Mitchell S. Problem-based learning: a review of literature on its outcomes and implementation issues. *Acad Med*. 1993/01/01. 1993;68(1):52–81. <https://doi.org/10.1097/00001888-199301000-00012>
12. Hakim A, Gibson R, Johansson L. Problem-based learning as a tool in postgraduate medical education. *Euro Resp J*. 2017 Sep 1;50(suppl 61):PA2770. <https://doi.org/10.1183/1393003.congress-2017.PA2770>.
13. Hurst JW. The overlecturing and underteaching of clinical medicine. *Arch Intern Med*. 2004;164(15):1605. <https://dx.doi.org/10.1001/archinte.164.15.1605>
14. Graffam B. Active learning in medical education: strategies for beginning implementation. *Med Teach*. 2007 Feb;29(1):38–42. <https://doi.org/10.1080/01421590601176398>.
15. Karunathilake I. Role of Problem-Based Learning (PBL) in postgraduate medical education. *J Postgraduate Inst Med*. 2019 Jul 31;6(1):91. <https://doi.org/10.4038/jpgim.8255>
16. Zhao W, He L, Deng W, Zhu J, Su A, Zhang Y. The effectiveness of the combined problem-based learning (PBL) and case-based learning (CBL) teaching method in the clinical practical teaching

- of thyroid disease. *BMC Med Educ.* 2020;20(1).
<https://dx.doi.org/10.1186/s12909-020-02306-y>
17. Albanese M. Problem-based learning: why curricula are likely to show little effect on knowledge and clinical skills. *Med Educ.* 2000 Sep;34(9):729–38. <https://doi.org/10.1046/j.1365-2923.2000.00753.x>
 18. Covidence. Veritas Health Innovation.
 19. Emily Jones M. Lib guides: creating a PRISMA flow diagram: new. PRISMA. 2020;
 20. Cook DA, Reed DA. Appraising the quality of medical education research methods: the medical education research study quality instrument and the Newcastle–Ottawa scale-education. *Acad Med.* 2015;90(8):1067–76.
<https://doi.org/10.1097/ACM.0000000000000786>
 21. StataCorp L. *Stata statistical software: Release 15* (2017). College Station, TX: StataCorp LP. 2017;
 22. Smidt A, Balandin S, Sigafos J, Reed VA. The Kirkpatrick model: A useful tool for evaluating training outcomes. *J Intellect Dev Disabil.* 2009;34(3):266–74.
<https://doi.org/10.1080/13668250903093125>
 23. Heading GS, Fuller JD, Lyle DM, Madden DL. Using problem-based learning in public health service based training. *N S W Public Health Bull.* 2007;18(1–2):8–13.
<https://doi.org/10.1071/NB07008>
 24. Sass P, Edelsack P. Teaching community health assessment skills in a problem-based format. *Acad Med.* 2001;76(1):88–91.
<https://doi.org/10.1097/00001888-200101000-00023>
 25. Cook DA, Thompson WG, Thomas KG, Thomas MR. Lack of interaction between sensing-intuitive learning styles and problem-first versus information-first instruction: a randomized crossover trial. *Adv Health Sci Ed.* 2009;14(1):79–90.
<https://doi.org/10.1007/s10459-007-9089-8>
 26. Foley RP, Levy J, Russinof HJ, Lemon MR. Planning and implementing a problem-based learning rotation for residents. *Teach Learn Med.* 1993;5(2):102–6.
<https://doi.org/10.1080/10401339309539600>
 27. Goodyear HM. Problem based learning in a junior doctor teaching programme. *Arch Dis Child.* 2005;90(3):275–8.
<https://doi.org/10.1136/adc.2003.037630>
 28. Ozuah PO, Curtis J, Stein REK. Impact of problem-based learning on residents' self-directed learning. 2001;155(June):669–73.
<https://doi.org/10.1001/archpedi.155.6.669>
 29. Heru AM. Teaching psychosomatic medicine using problem-based learning and role-playing. *Acad Psych.* 2011;35(4):245–8.
<https://doi.org/10.1176/appi.ap.35.4.245>
 30. Schultz-Ross RA, Kline AE. Using problem-based learning to teach forensic psychiatry. *Acad Psych.* 1999;23(1):37–41.
<https://doi.org/10.1007/BF03340034>
 31. Shefet D, Levkovitz Y. A national educational interactive conference for psychiatry residents in Israel. *Acad Psychiatry* 2011 Nov 1;35(6):419–20.
<https://doi.org/10.1176/appi.ap.35.6.419>
 32. Nguyen L, Brunnicardi FC, Dibardino DJ, Scott BG, Awad SS, Bush RL, et al. Education of the modern surgical resident: Novel approaches to learning in the era of the 80-hour workweek. *World J Surg.* 2006;30(6):1120–7.
<https://doi.org/10.1007/s00268-005-0038-5>
 33. Yates WR, Gerdes TT. Problem-Based Learning in Consultation. *Gen Hosp Psychiatry.* 1996;1638343(96):139–44.
[https://doi.org/10.1016/0163-8343\(96\)00022-9](https://doi.org/10.1016/0163-8343(96)00022-9)
 34. Prislun MD, Lenahan P, Shapiro J, Radecki S. Family practice residency behavioral science training: influence on graduate practice activity. *Fam Med.* 1997;29(7):483–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9232409>
 35. Lee L, Brunnicardi FC, Scott BG, et al. Impact of a novel education curriculum on surgical training within an academic training program. *J Surg Res.* 2008;145(2):308–12.
<https://doi.org/10.1016/j.jss.2007.03.046>
 36. Mulder SF, Bleijenberg G, Verhagen SC, Stuyt PMJ, Schijven MP, Tack CJ. Improved competence after a palliative care course for internal medicine residents. *Palliat Med.* 2009;23(4):360–8.
<https://doi.org/10.1177/0269216308100250>
 37. Carrero E, Gomar C, Penzo W, Rull M. Comparison between lecture-based approach and case/problem-based learning discussion for teaching pre-anaesthetic assessment. *Eur J Anaesthesiol.* 2007;24(12):1008–15.
<https://doi.org/10.1017/S0265021506002304>
 38. Komasa N, Berg BW, Minami T. Problem-based learning for anesthesia resident operating room crisis management training. *PLoS One.* 2018;13(11):1–10.
<https://doi.org/10.1371/journal.pone.0207594>
 39. Grol R, Mokkink H, Helsper-Lucas A, Tielens V, Bulte J. Effects of the vocational training of general practice consultation skills and medical performance. *Med Educ.* 1989;23(6):512–21.
<https://doi.org/10.1111/j.1365-2923.1989.tb01577.x>
 40. Rodrigues de Oliveira Filho G, Schonhorst L. Problem-based learning implementation in an intensive course of anaesthesiology: a preliminary report on residents' cognitive performance and perceptions of the educational environment. *Med Teach.* 2005;27(4):382–4.
<https://doi.org/10.1080/01421590500151021>
 41. Sakai T, Karausky PL, Valenti SL, Sandusky SL, Hirsch SC, Xu Y. Use of a problem-based learning discussion format to teach anaesthesiology residents research fundamentals. *J Clin Anesth.* 2013 Sep;25(6):434–8.
<https://doi.org/10.1016/j.jclinane.2013.04.008>
 42. Schmidt HG. Problem-based learning: rationale and description. *Med Educ.* 1983 Jan;17(1):11–6.
<https://doi.org/10.1111/j.1365-2923.1983.tb01086.x>
 43. Itani KMF, Miller CC, Church HM, McCollum CH. Impact of a problem-based learning conference on surgery residents' in training exam (ABSITE) scores. *J Surg Res.* 1997;70(1):66–8.
<https://doi.org/10.1006/jsre.1997.5112>
 44. Gruppen LD, Stansfield RB, Zhao Z, Sen S. Institution and specialty contribute to resident satisfaction with their learning environment and workload. *Acad Med.* 2015 Nov;90(11 Suppl):S77-82.
<https://doi.org/10.1097/ACM.0000000000000898>
 45. Thrush CR, Hicks EK, Tariq SG, et al. Optimal learning environments from the perspective of resident physicians and associations with accreditation length. *Acad Med.* 2007 Oct;82(10 Suppl):S121-5.
<https://doi.org/10.1097/ACM.0b013e318140658f>
 46. Cook DA, West CP. Perspective: Reconsidering the focus on “outcomes research” in medical education: a cautionary note.

Acad Med. 2013 Feb;88(2):162–7.

<https://doi.org/10.1097/ACM.0b013e31827c3d78>

47. Dweck C. Self-theories, their role in motivation, personality and development. Philadelphia, PA: Psychology Press; 2000.
48. Newman M, Van Den Bossche P, et al. Responses to the pilot systematic review of problem-based learning. *Med Educ.* 2004;38(9):921–3. <https://doi.org/10.1111/j.1365-2929.2004.01943.x>.
49. Verma AA, Pasricha S V, Jung HY, et al. Assessing the quality of clinical and administrative data extracted from hospitals: the General Medicine Inpatient Initiative (GEMINI) experience. *J Am Med Inform Assoc.* 2021 Mar 1;28(3):578–87. <https://doi.org/10.1093/jamia/ocaa225>
50. Gonzalez ML, Salmoni AJ. Online problem-based learning in postgraduate medical education - content analysis of reflection comments. *Teach Higher Educ.* 2008;13(2):183–92. <https://doi.org/10.1080/13562510801923302>.